

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Loamy (Ly) 5-9" Wind River Basin Precipitation Zone,

Site ID: R032XY222WY

Major Land Resource Area: 32 – Northern Intermountain Desertic Basins

Physiographic Features

This site occurs on nearly level to gently undulating rolling land and on slopes generally less than 20%.

Landform: Hillsides, alluvial fans, ridges & stream terraces

Aspect: N/A

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	4500	6600
Slope (percent):	0	30
Water Table Depth (inches):	None within 60 inches	
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):	0	0
Frequency:	None	None
Duration:	None	None
Runoff Class:	negligible	high

Climatic Features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the “Pavillion” climate station:

	<u>Minimum</u>	<u>Maximum</u>	<u>5 yrs. out of 10 between</u>
Frost-free period (days):	95	175	May 19 – September 19
Freeze-free period (days):	98	185	May 6 – October 3
Mean Annual Precipitation (inches):	2.50	12.54	

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53°F (30.5°F Avg. Min. to 58.5°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include “Riverton”, “Arminto”, and “Lost Cabin”.

Influencing Water Features

Wetland Description:	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
None	None	None	None	None

Stream Type: None

Representative Soil Features

The soils of this site are moderately deep to very deep (greater than 20" to bedrock), moderately well to well-drained & moderately slow to moderate permeable. The soil characteristic having the most influence on the plant community is the available moisture and the potential to develop soluble salts near the surface.

Major Soil Series correlated to this site include: Lostwells, Rairdent, Emblem, and Frisite.

Other Soil Series correlated to this site in MLRA 32 include:

Parent Material Kind: alluvium and residuum

Parent Material Origin: sandstone, shale

Surface Texture: loam, fine sandy loam, sandy loam

Surface Texture Modifier: none is most common but gravelly may occur

Subsurface Texture Group: loam, sandy clay loam, clay loam, silty clay, sandy loam

Surface Fragments ≤ 3" (% Cover): 0, occasionally up to 10

Surface Fragments > 3" (%Cover): 0

Subsurface Fragments ≤ 3" (% Volume): typically 0, occasionally up to 15

Subsurface Fragments > 3" (% Volume): typically 0, occasionally up to 10

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	moderately well drained	well drained
Permeability Class:	moderately slow	moderate
Depth (inches):	20	>60
Electrical Conductivity (mmhos/cm) ≤20":	0	8
Sodium Absorption Ratio ≤20":	0	13

Soil Reaction (1:1 Water) $\leq 20''$:	7.4	9.0
Soil Reaction (0.1M CaCl₂) $\leq 20''$:	NA	NA
Available Water Capacity (inches) $\leq 30''$:	3.0	6.3
Calcium Carbonate Equivalent (percent) $\leq 20''$:	0	20

Plant Communities

Ecological Dynamics of the Site:

Potential vegetation on this site is dominated by mid cool-season perennial grasses. Other significant vegetation includes winterfat, big sagebrush, and a variety of forbs. The expected potential composition for this site is about 75% grasses, 10% forbs and 15% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.


As this site deteriorates species such as blue grama, Sandberg bluegrass, and big sagebrush will increase. Weedy annuals will invade. Cool-season grasses such as, rhizomatous wheatgrasses, needleandthread, and Indian ricegrass will decrease in frequency and production.

Big sagebrush may become dominant on areas with an absence of fire and sufficient amount of precipitation. Wildfires are actively controlled in recent times and as a result old decadent stands of big sagebrush persist. Chemical control using herbicides has replaced the historic role of fire on this site. Recently, prescribed burning has regained some popularity.

Due to the amount and pattern of the precipitation, the big sagebrush component may not be resilient once it has been removed or severely reduced if a vigorous stand of grass exists and is maintained. On these areas, blue grama may become dominant if the area is subjected to a combination of frequent and severe grazing, especially yearlong grazing. As a result, a dense sod cover of blue grama will become established.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



- BM** - Brush Management (fire, chemical, mechanical)
- Freq. & Severe Grazing** - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season
- GLMT** - Grazing Land Mechanical Treatment
- LTPG** - Long-term Prescribed Grazing
- MCSLG** - Moderate, Continuous Season-long Grazing
- NU, NF** - No Use and No Fire
- PG** - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)
- VLTPG** - Very Long-term Prescribed Grazing (could possibly take generations)
- WF** - Wildfire (Natural or Human Caused)

Plant Community Composition and Group Annual Production
Reference Plant Community (HCPC)

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Annual Production (Normal Year)		
			Group	lbs./acre	% Comp.
GRASSES AND GRASS-LIKES					
GRASSES/GRASSLIKES					
Western wheatgrass or	Pascopyrum smithii	PASM	1	80 - 160	20 - 40
Thickspike wheatgrass	Elymus lanceolatus ssp. lanceolatus	ELLAL	1	80 - 160	20 - 40
Griffiths wheatgrass or	Elymus albicans	ELAL7	2	20 - 40	5 - 10
Bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6			
Needleandthread grass	Hesperostipa comata	HECO26	3	40 - 80	10 - 20
Indian ricegrass	Achnatherum hymenoides	ACHY	4	20 - 60	5 - 15
Bottlebrush squirreltail	Elymus elymoides	ELELE	5	20 - 40	5 - 10
MISC. GRASSES/GRASSLIKES			6	20 - 60	5 - 15
Prairie junegrass	Koeleria macrantha	KOMA	6	0 - 20	0 - 5
Sandberg bluegrass	Poa secunda	POSE	6	0 - 20	0 - 5
Threadleaf sedge	Carex filifolia	CAFI	6	0 - 20	0 - 5
Blue grama	Bouteloua gracilis	BOGR2	6	0 - 20	0 - 5
Fendler threeawn	Aristida purpurea var. longiseta	ARPUL	6	0 - 20	0 - 5
other perennial grasses (native)		2GP	6	0 - 20	0 - 5
FORBS			7	20 - 40	5 - 10
Scarlet globemallow	Sphaeralcea coccinea	SPCO	7	0 - 20	0 - 5
Hood's phlox	Phlox hoodii	PHHO	7	0 - 20	0 - 5
Wild onion	Allium textile	ALTE	7	0 - 20	0 - 5
Fringed sagewort	Artemisia frigida	ARFR4	7	0 - 20	0 - 5
Parsnipflower buckwheat	Eriogonum heracleoides	ERHE2	7	0 - 20	0 - 5
Missouri milkvetch	Astragalus missouriensis	ASMI10	7	0 - 20	0 - 5
White loco	Oxytropis sericea	OXSES2	7	0 - 20	0 - 5
Leafy wildparsley	Musineon divaricatum	MUDI	7	0 - 20	0 - 5
Toadflax	Comandra umbellata	COUM	7	0 - 20	0 - 5
Wavyleaf paintbrush	Castilleja applegatei martinii	CAAPM	7	0 - 20	0 - 5
Threadleaf fleabane	Erigeron filifolius	ERFI2	7	0 - 20	0 - 5
Smooth woodyaster	Xylorhiza glabruiscula	XUGL	7	0 - 20	0 - 5
Bigseed biscuitroot	Lomatium macrocarpum	LOMA3	7	0 - 20	0 - 5
Tapertip hawksbread	Crepis acuminata	CRAC2	7	0 - 20	0 - 5
Small-leaf pussytoes	Antennaria parvifolia	ANPA4	7	0 - 20	0 - 5
Little larkspur	Delphinium bicolor	DEBI	7	0 - 20	0 - 5
Meadow deathcamas	Zigadenas venenosus	ZIVE	7	0 - 20	0 - 5
Plains pricklypear cactus	Opuntia polyacantha	OPPO	7	0 - 20	0 - 5
other perennial forbs (native)		2FP	7	0 - 20	0 - 5
TREES/SHRUBS					
Wyoming big sagebrush	Artemisia tridentata wyomingensis	ARTRW8	8	20 - 60	5 - 15
Winterfat	Krascheninnikovia lanata	KRAL2	9	0 - 20	0 - 5
MISC. SHRUBS			10	20 - 40	5 - 10
Rubber rabbitbrush	Ericameria nauseosa	ERNA10	10	0 - 20	0 - 5
Green rabbitbrush	Chrysothamnus viscidiflorus	CHVI8	10	0 - 20	0 - 5
Spiny gilia	Leptodactylon pungens	LEPU	10	0 - 20	0 - 5
other shrubs & half shrubs (native)		2SHRUB	10	0 - 20	0 - 5

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

Rhizomatous Wheatgrasses/Needleandthread Plant Community

This plant community is the interpretive plant community for this site and is considered to be the Historic Climax Plant Community (HCPC). This state evolved with grazing by large herbivores and periodic fires. The cyclical nature of the fire regime in this community prevented big sagebrush from being the dominant landscape. This plant community can be found on areas that are properly managed with grazing and/or prescribed burning, and on areas receiving occasional short periods of rest. The potential vegetation is about 75% grasses or grass-like plants, 10% forbs, and 15% woody plants. This state is dominated by cool season mid-grasses.

The major grasses include western wheatgrass, needleandthread, and Indian ricegrass. Other grasses occurring in this state include thickspike wheatgrass, bluebunch wheatgrass, and bottlebrush squirreltail. A variety of forbs and half-shrubs also occur, as shown in the preceding table. Big sagebrush is a conspicuous element of this state, occurs in a mosaic pattern, and makes up 5 to 15% of the annual production. Winterfat is a common component found on this site.

The total annual production (air-dry weight) of this state is about 400 lbs./acre, but it can range from about 225 lbs./acre in unfavorable years to about 600 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This plant community is extremely stable and well adapted to the Northern Intermountain Desertic Basins climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Moderate, continuous season-long grazing will convert the plant community to the *Perennial Grass/Big Sagebrush Plant Community*. Prolonged Drought will exacerbate this transition.

Perennial Grass/Big Sagebrush Plant Community

Historically, this plant community evolved under grazing by large ungulates and a low fire frequency. Currently, it is found under moderate, season-long grazing by livestock and will be exacerbated by

prolonged drought conditions. In addition, the fire regime for this site has been modified and extended periods without fire is now common. This plant community is still dominated by cool-season grasses, while short warm-season grasses and miscellaneous forbs account for the balance of the understory. Wyoming big sagebrush is now a conspicuous part of the overall production and accounts for the majority of the overstory.

Dominant grasses include western wheatgrass, and needleandthread. Grasses and grass-like species of secondary importance include blue grama, Sandberg bluegrass and threadleaf sedge. Forbs commonly found in this plant community include scarlet globemallow, fringed sagewort, hairy goldaster, and phlox. Sagebrush can make up to 25% of the annual production. The overstory of sagebrush and understory of grass and forbs provide a diverse plant community.

When compared to the Historic Climax Plant Community, big sagebrush and blue grama have increased. Plains pricklypear cactus will also have increased, but occurs only in small patches. Indian ricegrass has decreased and may occur in only trace amounts under the sagebrush canopy or within the patches of pricklypear. In addition, the amount of winterfat may or may not have changed depending on the season of use.

The total annual production (air-dry weight) of this state is about 320 pounds per acre, but it can range from about 180 lbs./acre in unfavorable years to about 480 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. The herbaceous component is mostly intact and plant vigor and replacement capabilities are sufficient. Water flow patterns and litter movement may be occurring but only on steeper slopes. Incidence of pedestalling is minimal. Soils are mostly stable and the surface shows minimum soil loss. The watershed is functioning and the biotic community is intact.

Transitions or pathways leading to other plant communities are as follows:

- Prescribed grazing or possibly long-term prescribed grazing, will convert this plant community to the *HCPC*. The probability of this occurring is high especially if rotational grazing along with short deferred grazing is implemented as part of prescribed method of use. In addition, the removal of fire suppression will allow a somewhat natural fire regime to reoccur to more easily transition between this plant community and the *HCPC*. A prescribed fire treatment can be useful to hasten this transition, if desired.
- Frequent and severe grazing plus no fire, will convert the plant community to the Big *Sagebrush/Bare Ground Plant Community*. The probability of this occurring is high. This is especially evident on areas with historically higher precipitation and the sagebrush stand is not adversely impacted by drought or heavy browsing.

- Frequent and severe grazing (yearlong grazing) plus wildfire or brush control, will convert the plant community to the *Blue Grama Sod Plant Community*. The probability of this occurring is high, especially if the sagebrush stand has been severely affected by drought or heavy use or has been removed altogether.
- Frequent and severe grazing (yearlong grazing) on more saline soils, will convert the plant community to the *Salt Tolerant Shrub/Bare Ground Plant Community*. The probability of this occurring is high especially on soils with elevated salts and the sagebrush stand has been severely affected by drought and heavy use or has been removed altogether.

Big Sagebrush/Bare Ground Community

This plant community is the result of frequent and severe grazing and protection from fire. Sagebrush dominates this plant community, as the annual production of sagebrush exceeds 25%. Wyoming big sagebrush is a significant component of the plant community and the preferred cool season grasses have been eliminated or greatly reduced.

The dominant grasses are Sandberg bluegrass, and blue grama. Weedy annual species such as cheatgrass may occupy the site if a seed source is available. Cactus and sageworts often increase. Noxious weeds such as Russian Knapweed, leafy spurge, or Canada thistle may invade the site if a seed source is available. The interspaces between plants have expanded leaving the amount of bare ground more prevalent. As compared with the HCPC or the Perennial Grass/Big Sagebrush Plant Communities, the annual production is similar, as the shrub production compensates for the decline in the herbaceous production.

The total annual production (air-dry weight) of this state is about 300 pounds per acre, but it can range from about 200 lbs./acre in unfavorable years to about 400 lbs./acre in above average years.

The following is the growth curve of the plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change as the stand becomes more decadent. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the sagebrush plants is increased. Continued frequent and severe grazing or the removal of grazing does not seem to affect the composition or structure of the plant community. Plant diversity is moderate to poor. The plant vigor is diminished and replacement capabilities are limited due to the reduced number of cool-season grasses. Plant litter is noticeably less when compared to the HCPC.

Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels may be noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope.

Transitions or pathways leading to other plant communities are as follows:

- Brush management, followed by prescribed grazing, will return this plant community at or near the HCPC. If prescribed fire is used as a means to reduce or remove the shrubs, sufficient fine fuels will need to be present. This may require deferment from grazing prior to treatment. Post management is critical to ensure success. This can range from two or more years of rest to partial growing season deferment, depending on the condition of the understory at the time of treatment and the growing conditions following treatment. In the case of an intense wildfire that occurs when desirable plants are not completely dormant, the length of time required to reach the HCPC may be increased and seeding of natives are recommended.
- Brush management, followed by frequent and severe grazing, will convert the plant community to the *Blue Grama Sod Plant Community*.

Blue Grama Sod Plant Community

This plant community is the result of frequent and severe yearlong grazing, which have adversely affected the perennial grasses as well as the addition of other impacts that can affect the shrub component. These factors include drought and wildfires, and human brush control measures. A dense sod of blue grama with patches of threadleaf sedge dominates this state. Pricklypear cactus can become dense enough in patches so that livestock cannot graze forage growing within the cactus clumps. Big sagebrush has been reduced to small patches or in some cases removed.

When compared to the Historic Climax Plant Community, blue grama, threadleaf sedge, and pricklypear have increased. All cool-season mid-grasses, forbs, and most shrubs have been greatly reduced. Production has been significantly decreased.

The total annual production (air-dry weight) of this state is about 100 pounds per acre, but it can range from about 55 lbs./acre in unfavorable years to about 150 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This sod is extremely resistant to change and continued frequent and severe grazing or the removal of grazing does not seem to affect the plant composition or structure of the plant community. The biotic integrity of this state is not functional and plant diversity is extremely low. The plant vigor is significantly weakened and replacement capabilities are limited due to the reduced number of cool-season grasses.

This sod bound plant community is very resistant to water infiltration. While this sod protects the site itself, off-site areas are affected by excessive runoff that can cause rills and gully erosion. Water flow patterns are obvious in the bare ground areas and pedestalling is apparent along the sod edges. Rill channels are noticeable in the interspaces and down slope. The watershed may or may not be functioning, as runoff may affect adjoining sites.

Transitions or pathways leading to other plant communities are as follows:

- Grazing land mechanical treatment (chiseling, etc.) and pricklypear cactus control (if needed), followed by prescribed grazing, will return this plant community to near *Historic Climax Plant*

Community condition. Seeding with native species may also be needed depending on how far a shift away from mid cool-season grasses has occurred.

Salt Tolerant Shrub/Bare Ground Plant Community

This plant community can occur where states are subjected to continuous yearlong grazing and where soils are influenced by elevated amounts of soluble salts. Salt tolerant shrubs are a significant component of the plant community and the preferred cool season grasses have been eliminated or greatly reduced. Wyoming big sagebrush is still a component of the plant community but may not be the primary overstory species.

This site is dominated by an overstory of shrubs, which can vary widely in the composition and production. This variation results from the dissimilarity in quantity of soluble salts present in the soils and the availability of shrubs to occupy the site. The dominant shrubs are greasewood, big sagebrush, rubber rabbitbrush, and a number of different saltbushes.

Perennial cool season mid-grasses have been reduced or removed, leaving mostly patches of blue grama and annuals. Cheatgrass and weedy annual forbs such as halogeton, Russian thistle, and kochia will occupy the site if a seed source is available. Noxious weeds such as Russian Knapweed may also invade this site.

The interspaces between plants have expanded leaving the amount of bare ground more prevalent. Surface salts have increased, especially on sites dominated by greasewood and saltbushes. The leaves of these plants contain high amounts of sodium and other salts, and when shed these soluble salts are transferred to the soils underneath the plants. Consequently, the soil can exhibit wide variations in soluble salts, which can explain the variation in shrub composition.

The total annual production (air-dry weight) of this state is about 225 pounds per acre, but it can range from about 125 lbs./acre in unfavorable years to about 350 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change. These areas are actually more resistant to fire as less fine fuels are available and the bare ground between the shrubs has increased. Continued frequent and severe grazing does not affect the composition or structure of the plant community. Plant diversity is moderate to poor. The biotic integrity of this state is mostly dysfunctional because of the predominant salt tolerant shrub overstory and absence of perennial cool-season grasses.

Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels may be noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope.

Transitions or pathways leading to other plant communities are as follows:

- Prescribed grazing or possibly long-term prescribed grazing, will convert this plant community to the *Salt Tolerant Shrub/Rhizomatous Wheatgrass Vegetative State*. Recovery to near *Historic Climax Plant Community* condition is difficult to impossible due to the resistance of

these shrubs to herbicides and other brush management techniques. In addition, the increase in surface salts has had accumulated effects on the soil so most of the herbaceous plants associated with the HCPC are no longer suitable for this site. The most notable exception is the rhizomatous wheatgrasses and bottlebrush squirreltail. Soil remediation to reduce the surface salts is not recommended, as this is mostly ineffective and extremely costly. Seeding more salt-tolerant native grasses and forbs will improve plant cover and the productivity of the site.

Salt Tolerant Shrub/Rhizomatous Wheatgrasses Plant Community

This plant community can occur where the Salt Tolerant Shrub/Bare Ground Plant Community is rested and a prescribed grazing management practice is implemented. Salt tolerant shrubs and Wyoming big sagebrush remain a significant component of the plant community but preferred cool season grasses have reestablished.

This site is dominated by an overstory of a variety of shrubs, such as Wyoming big sagebrush, rubber rabbitbrush, greasewood, and a variety of saltbushes. Some perennial cool season mid-grasses have once again reestablished such as rhizomatous wheatgrasses and bottlebrush squirreltail. Other important grasses include Sandberg bluegrass and blue grama. Patches of annuals such as cheatgrass and weedy annual forbs such as halogeton, Russian thistle, and kochia will persist on this site. Noxious weeds such as Russian Knapweed may also remain if not treated. The interspaces between plants will have diminished in size. When compared with the HCPC or the Perennial Grass/Big Sagebrush Plant Communities, the annual production is similar, but the plant species are clearly unique.

The total annual production (air-dry weight) of this state is about 300 pounds per acre, but it can range from about 200 lbs./acre in unfavorable years to about 425 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	50	25	5	0	10	0	0	0

(Monthly percentages of total annual growth)

This plant community is mostly resistant to change, but species composition can be altered through long-term overgrazing. The herbaceous component is stable, but does not include most historical species. Plant vigor and replacement capabilities are sufficient. The biotic community is not intact because of the predominant salt tolerant shrub overstory and lack of historic grass species. Plant diversity is moderate.

Soils are mostly stable and recent soil loss is minimal. This should not be confused with evidence of remnant erosion. Water flow patterns and litter movement is stable but is still occurring on steeper slopes. Incidence of pedestalling is improving. The watershed may or may not be functioning.

Transitions or pathways leading to other plant communities are as follows:

- Frequent and severe grazing will convert the plant community to the *Salt Tolerant Shrub/Bare Ground Plant Community*.
- Recovery to near *Historic Climax Plant Community* condition is difficult to impossible due to the resistance of these shrubs to herbicides and other brush management techniques. In

addition, the increase in surface salts has had accumulated effects on the soil so most of the herbaceous plants associated with the HCPC are no longer suitable for this site. The most notable exception is the rhizomatous wheatgrasses and bottlebrush squirreltail. Soil remediation to reduce the surface salts is not recommended, as this is mostly ineffective and extremely costly. Seeding more salt-tolerant grasses and forbs will improve the productivity of site and plant cover, but will not improve the biotic integrity.

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

Rhizomatous Wheatgrasses/Needleandthread (HCPC): The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

Perennial Grass/Big Sagebrush Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants and hosts of other nesting birds utilize stands in the 20-30% cover range.

Big Sagebrush/Bare Ground Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15% protein and 40-60% digestibility during that time. This community provides nesting habitat for sage grouse. However, it provides little foraging opportunities for upland game birds, as fewer forbs are available. Many grassland obligate small mammals would occur here.

Blue Grama Sod Plant Community: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Perennial Grass/ Big Sagebrush Plant Community is limited. Generally, these are not target plant communities for wildlife habitat management.

Salt Tolerant Shrub/Bare Ground Plant Community: This plant community exhibits a low level of plant species diversity due to the accumulation of salts near the soil surface. It does, however, provide thermal and escape cover for upland birds. Upland game birds may find little foraging opportunities, as fewer forbs are available. Many grassland obligate small mammals would occur here.

Salt Tolerant Shrub/Rhizomatous Wheatgrass Plant Community: This plant community can provide winter foraging for elk, mule deer and antelope. This community provides nesting habitat for upland game birds. However, it provides little foraging opportunities for upland game birds, as fewer forbs are available. Many grassland obligate small mammals would occur here.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 32XY, 5-9 inch Wind River Basin

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope
GRASSES/GRASSLIKE							
alkali bluegrass	<i>Poa secunda</i> ssp. <i>juncifolia</i>	POSEJ	DDDD	PPPP	DDDD	PPPP	PPPP
alkali cordgrass	<i>Spartina gracilis</i>	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU
alkali sacaton	<i>Sporobolus airoides</i>	SPAI	PPPP	DDDD	PPPP	DDDD	DDDD
American manna grass	<i>Glyceria grandis</i>	GLGR	DDDD	UUUU	DDDD	UUUU	UUUU
American sloughgrass	<i>Beckmannia syzigachne</i>	BESY	DDDD	UUUU	DDDD	UUUU	UUUU
Baltic rush	<i>Juncus balticus</i>	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU
basin wildrye	<i>Leymus cinereus</i>	LEC4	PPPP	PPPP	PPPP	DDDD	DDDD
beaked sedge	<i>Carex rostrata</i>	CARO6	DDDD	UUUU	DDDD	UUUU	UUUU
bearded wheatgrass	<i>Elymus caninus</i>	ELCA	PPPP	DDDD	PPPP	DDDD	DDDD
big bluegrass	<i>Poa ampla</i> (syn. to <i>Poa secunda</i>)	POAM (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
blue grama	<i>Bouteloua gracilis</i>	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD
bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD
bottlebrush squirreltail	<i>Elymus elymoides</i>	ELELE	DDDD	DDDD	DDDD	UUUU	UUUU
bulrush	<i>Scirpus</i> spp.	SCIRP	DDDD	UUUU	DDDD	UUUU	UUUU
Canada wildrye	<i>Elymus canadensis</i>	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD
Fendler threeawn	<i>Aristida purpurea longiseta</i>	ARFUL	UUUU	UUUU	UUUU	UUUU	UUUU
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP
inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU
little bluestem	<i>Schizachyrium scoparium</i>	SCSC	PPPP	PPPP	PPPP	DDDD	DDDD
mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	<i>Carex nebrascensis</i>	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD
needleandthread	<i>Hesperostipa comata</i>	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP
northern reedgrass	<i>Calamagrostis stricta</i>	CAST13	PPPP	DDDD	PPPP	UUUU	UUUU
Nuttall's alkali grass	<i>Puccinellia nuttalliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP
plains reedgrass	<i>Calamagrostis montanensis</i>	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD
prairie cordgrass	<i>Spartina pectinata</i>	SPPE	PPPP	DDDD	PPPP	UUUU	UUUU
prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD
prairie sandreed	<i>Calamovilfa longifolia</i>	CALO	PPPP	DDDD	PPPP	UUUU	UUUU
reed canarygrass	<i>Phalaris arundinacea</i>	PHAR3	DDDD	UUUU	DDDD	UUUU	UUUU
rush	<i>Juncus</i> spp.	JUNCU	DDDD	UUUU	DDDD	UUUU	UUUU
sand dropseed	<i>Sporobolus cryptandrus</i>	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU
Sandberg bluegrass	<i>Poa secunda</i>	POSE	DDDD	DDDD	DDDD	DDDD	DDDD
slender wheatgrass	<i>Elymus trachycaulis</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD
spike sedge	<i>Carex nardina</i>	CANA2	DDDD	DDDD	DDDD	UUUU	UUUU
thickspike wheatgrass	<i>Elymus lanceolatus</i>	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD
threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP
tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD
water sedge	<i>Carex aquatilis</i>	CAAQ	DDDD	UUUU	DDDD	UUUU	UUUU
western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DDDD	DDDD	DDDD	DDDD	DDDD
FORBS							
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU
American vetch	<i>Vicia americana</i>	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP
arrowgrass	<i>Triglochin</i> spp.	TRIGL	T	T	T	T	T
asters	<i>Aster</i> spp.	ASTER	UUUU	UUUU	UUUU	UUUU	UUUU
badlands mule-ears	<i>Wyethia scabra</i>	WYSC	UUUU	UUUU	UUUU	UUUU	UUUU
beaked skeletonweed	<i>Shinnersoseris rostrata</i>	SHRO2	UUUU	UUUU	UUUU	UUUU	UUUU
biscuitroots	<i>Lomatium</i> spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD
blue-eyed grass	<i>Sisyrinchium</i> spp.	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD
breadroot scurfpea	<i>Pediomelum esculentum</i>	PEES	DDDD	DDDD	DDDD	DDDD	DDDD
buttercandle	<i>Cryptantha celosiodes</i>	CRCE	UUUU	UUUU	UUUU	UUUU	UUUU
cattail, broad-leaf	<i>Typha latifolia</i>	TYLA	DDDD	UUUU	DDDD	UUUU	UUUU
cattail, narrow-leaf	<i>Typha angustifolia</i>	TYAN	DDDD	UUUU	DDDD	UUUU	UUUU
desert princeplume	<i>Stanleya pinnata</i>	STPIP	T	T	T	T	T
Douglas' dusty maid	<i>Chaenactis douglasii</i>	CHDO	UUUU	UUUU	UUUU	UUUU	UUUU
fleabane	<i>Erigeron</i> spp.	ERIGU	UUUU	UUUU	UUUU	UUUU	UUUU
foothills deathcamas	<i>Zigadenus paniculatus</i>	ZIPA2	T	T	T	T	T
fringed sagewort	<i>Artemisia frigida</i>	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU
green sagewort	<i>Artemisia dracunculoides</i>	ARDR4	UUUU	UUUU	UUUU	UUUU	UUUU
hawkbeard	<i>Crepis acuminata</i>	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD
horsetails	<i>Equisetum</i> spp.	EQUIS	UUUU	UUUU	UUUU	UUUU	UUUU
Indian paintbrush	<i>Castilleja</i> spp.	CASTI2	DDDD	DDDD	DDDD	DDDD	DDDD
iris	<i>Iris</i> spp.	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU
larkspur	<i>Delphinium</i> spp.	DELPH	DDDD	DDDD	DDDD	DDDD	DDDD
licorice-root	<i>Ligusticum</i> spp.	LIGUS	UUUU	UUUU	UUUU	UUUU	UUUU
lupine	<i>Lupinus</i> spp.	LUPIN	DDDD	T	DDDD	DDDD	DDDD
milkvetch	<i>Astragalus</i> spp.	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD
miner's candle	<i>Cryptantha virgata</i>	CRV14	UUUU	UUUU	UUUU	UUUU	UUUU
mustard	<i>Brassica</i> spp.	BRASS2	UUUU	UUUU	UUUU	UUUU	UUUU
nailwort	<i>Paronychia</i> spp.	PARON	UUUU	UUUU	UUUU	UUUU	UUUU
Nuttall's povertyweed	<i>Monolepis nuttalliana</i>	MONU	UUUU	UUUU	UUUU	UUUU	UUUU
penstemon	<i>Penstemon</i> spp.	PENST	PPPP	PPPP	PPPP	PPPP	PPPP
phlox	<i>Phlox</i> spp.	PHLOX	UUUU	UUUU	UUUU	UUUU	UUUU
plains springparsley	<i>Cymopterus acaulis</i>	CYAC	UUUU	DDDD	UUUU	UUUU	UUUU
poison hemlock	<i>Conium maculatum</i>	COMA2	T	T	T	T	T
prairie bluebells	<i>Mertensia lanceolata</i>	MELA3	DDDD	PPPP	DDDD	DDDD	DDDD
Pursh seepweed	<i>Suaeda calceoliformis</i>	SUCA2	UUUU	UUUU	UUUU	UUUU	UUUU
rosy pussytoes	<i>Antennaria rosea</i>	ANRO2	UUUU	UUUU	UUUU	UUUU	UUUU
sandwort	<i>Arenaria</i> spp.	ARENA	UUUU	UUUU	UUUU	UUUU	UUUU
silverweed cinquefoil	<i>Argentina anserina</i>	ARAN7	UUUU	UUUU	UUUU	UUUU	UUUU
stemless goldenweed	<i>Haplopappus acaulis</i>	HAAC	UUUU	UUUU	UUUU	UUUU	UUUU
sulphur flower buckwheat	<i>Eriogonum umbellatum</i>	ERUM	UUUU	UUUU	UUUU	UUUU	UUUU
tufted evening-primrose	<i>Oenothera caespitosa</i>	OECA10	UUUU	UUUU	UUUU	UUUU	UUUU
twogrooved milkvetch	<i>Astragalus bisulcatus</i>	ASB12	T	T	T	T	T
water hemlocks	<i>Cicuta</i> spp.	CICUT	T	T	T	T	T
western buttercup	<i>Ranunculus occidentalis</i>	ROAOC	DDDD	DDDD	DDDD	DDDD	DDDD
western dock	<i>Rumex aquaticus</i>	RUAQ	UUUU	UUUU	UUUU	UUUU	UUUU
western yarrow	<i>Achillea lanulosa</i>	ACHIL	UUUU	UUUU	UUUU	UUUU	UUUU
wild onion	<i>Allium textile</i>	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD
woodyaster	<i>Xylorhiza</i> spp.	XYLOR	T	T	T	T	T
woolly plantain	<i>Plantago patagonica</i>	PLPA2	UUUU	UUUU	UUUU	UUUU	UUUU

TREES, SHRUBS & HALF-SHRUBS							
big sagebrush	Artemisia tridentata	ARTR2	UUUU	DDDD	UUUU	DDDD	DDDD
birdfoot sagebrush	Artemisia pedatifida	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU
black greasewood	Sarcobatus vermiculatus	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD
black sagebrush	Artemisia nova	ARNO4	DDDD	PPPP	UUUU	PPPP	PPPP
broom snakeweed	Gutierrezia sarothrae	GUSA2	UUUU	UUUU	UUUU	UUUU	UUUU
bud sagebrush	Picrothamnus desertorum	PIDE4	PPPP	PPPP	DDDD	PPPP	PPPP
fourwing saltbush	Atriplex canescens	ATCA2	PPPP	PPPP	PPPP	PPPP	PPPP
Gardners saltbush	Atriplex gardneri	ATGA	PPPP	PPPP	DDDD	PPPP	PPPP
green rabbitbrush	Chrysothamnus viscidiflorous	CHV18	DDDD	DDDD	DDDD	DDDD	DDDD
plains cottonwood (sprouts)	Populus deltoides	PODEM	DDDD	DDDD	DDDD	DDDD	DDDD
Rocky Mountain juniper	Juniperus scopulorum	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU
rubber rabbitbrush	Ericameria nauseosa	ERNA10	UUUU	DDDD	UUUU	DDDD	DDDD
shadscale saltbush	Atriplex confertifolia	ATCO	UUUU	UUUU	UUUU	UUUU	UUUU
shortspine horsebrush	Tetradymia spinosa	TESP2	UUUU	UUUU	UUUU	UUUU	UUUU
silver sagebrush	Artemisia cana	ARCAC5	DDDD	DDDD	DDDD	PPPP	PPPP
silverberry	Eleagnus commutata	ELCO	UUUU	UUUU	UUUU	DDDD	UUUU
skunkbush sumac	Rhus trilobata	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD
spiny hopsage	Grayia spinosa	GRSP	UUUU	UUUU	UUUU	UUUU	UUUU
Utah juniper	Juniperus osteosperma	JUOS	UUUU	UUUU	UUUU	DDDD	UUUU
wax currant	Ribes cereum	RICE	UUUU	UUUU	UUUU	DDDD	DDDD
western snowberry	Symphoricarpos occidentalis	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU
wildrose	Rosa woodsii var. woodsii	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD
willows	Salix spp.	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU
winterfat	Krascheninnikovia lanata	KRLA2	PPPP	PPPP	PPPP	PPPP	PPPP
yucca	Yucca glauca	YUGL	DDDD	DDDD	DDDD	DDDD	DDDD

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (lb./ac)	Carrying Capacity* (AUM/ac)
Rhizomatous WG/Needleandthread	225-600	.20
Perennial Grass/Big Sagebrush	180-480	.16
Big Sagebrush/Bare Ground	200-400	.10
Blue Grama Sod	55-150	.05
Salt Tolerant Shrub/Bare Ground	125-350	.07
Salt Tolerant Shrub/Rhizomatous Wheatgrasses	200-425	.16

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderately slow to moderate. Runoff potential for this site varies from low to moderate depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational Uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood Products

No appreciable wood products are present on the site.

Other Products

None noted.

Supporting Information

Associated Sites

Shallow Loamy	R032XY262WY
Sandy	R032XY250WY
Clayey	R032XY204WY
Lowland	R032XY228WY

Similar Sites

() – Loamy 10-14" Foothills and Basins East P.Z., R032XY322WY has higher production than Loamy 5-9" WR.

Inventory Data References (narrative)

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everett Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties.

Inventory Data References

Ocular field estimations observed by trained personnel.

Site Correlation

This site occurs entirely within Wyoming.

Type Locality

Field Offices

Casper, Lander, Riverton, Fort Washakie, Dubois

Relationship to Other Established Classifications

Other References

Site Description Approval

State Range Management Specialist

Date